

The Malaysian Consensus on
the Management of Acute and Persistent Hyperkalaemia: A Multidisciplinary Approach
2024

Placeholder for MSN	NHAM	College of Emergency Physicians
---------------------	------	---------------------------------

DEVELOPMENT COMMITTEE

Chairpersons

Professor Dr Lim Soo Kun
Consultant Nephrologist
University Malaya Medical Centre

Dr Mohd Rahal Yusoff
Consultant Cardiologist,
Hospital Universiti Teknologi MARA

Professor Dr Shaik Farid Abdul Wahab
Consultant Emergency Medicine Physician,
Hospital Universiti Sains Malaysia, Kubang Kerian

Development Committee

Consultant Cardiologists

Dato' Dr Azmee Mohd Ghazi
Institut Jantung Negara

Dato' David Chew Soon Ping
CVS Kuala Lumpur

Dr Liew Hounng Bang
Hospital Queen Elizabeth, Kota Kinabalu

Consultant Emergency Medicine Physicians

Dr Paranthaman Kaneson
Subang Jaya Medical Centre

Dr Siti Suhaila Hamzah
Hospital Sungai Buloh

Consultant Nephrologists

Dr Ching Chen Hua
Hospital Sultanah Bahiyah, Alor Setar

Dr Prasad Menon
Subang Jaya Medical Centre

Dr Sunita Bavanandan
Hospital Kuala Lumpur

Dr Tan Li Ping
Ara Damansara Medical Centre

SUMMARY

Hyperkalaemia (HyperK⁺) is a potentially life-threatening condition that poses significant challenges in both acute and chronic care, especially for patients with chronic kidney disease, heart failure, and diabetes mellitus. While renin-angiotensin-aldosterone system inhibitors (RAASi) are a guideline-directed medical therapy known for their cardiorenal benefits, concerns about the increased risk of hyperK⁺ associated with these medications have hindered their optimal use. Despite these challenges, ambiguity remains regarding the management of hyperK⁺, highlighting the need for a standardised approach. This Malaysian consensus provides evidence-based guidelines for managing acute and persistent hyperK⁺ tailored to the local healthcare context.

A multidisciplinary panel of experts, including nephrologists, cardiologists, and emergency physicians, reviewed current evidence and international guidelines to develop 16 consensus statements. These statements address key areas, which include the definition of hyperK⁺, emergency management strategies, long-term treatment with novel potassium binders, and collaborative care practices. The consensus emphasises early identification, individualised care, and the role of novel therapies like sodium zirconium cyclosilicate and patiromer in optimising RAASi therapy.

Through this consensus, the panel aims to improve outcomes for cardiorenal patients by providing a structured approach to managing hyperK⁺ in Malaysia.

CONSENSUS STATEMENTS DEVELOPMENT

The development of the consensus statements involved a comprehensive and structured process. A steering committee (SC) comprised of 12 members, including emergency physicians (3), nephrologists (4), and cardiologists (4), was formed to develop these statements.

The initial phase involved a scoping review of recent consensus statements and guidelines followed by the creation of 49 statements across six major themes, which were the definition of hyperK⁺, diagnosing and managing hyperK⁺ in emergency settings, preventing hyperK⁺ in at-risk cardiorenal patients, assessing risk and managing hyperK⁺ in cardiorenal patients, K⁺-lowering therapies for hyperK⁺ in cardiorenal patients, and collaborative care. These statements aimed to recommend diagnosis and treatment approaches for hyperK⁺ in both acute (emergency) and persistent settings. The initial statements were presented to the SC for extensive discussion. Through this process, the number of statements was reduced to 20, while maintaining the original six themes. The revised 20 statements were then developed into a survey using Microsoft Forms. All submissions were anonymised, and consent was implied when specialists answered the survey on a volunteer basis.

The survey was distributed to all registered specialists through their respective professional societies: the College of Emergency Physicians, the Malaysian Society of Nephrology, and the National Heart Association of Malaysia. Based on the combined number of registered specialists, the SC agreed that a response rate of approximately

30% would be acceptable. The survey duration was set for two months to allow optimal response time.

The statements were evaluated using a 4-point Likert scale, “Strongly agree, Agree, Disagree and Strongly disagree”, with an option to provide comments for each statement. Votes of “Strongly agree” and “Agree” were totalled to determine the level of agreement. Statements with $\geq 75\%$ agreement were included in the consensus.

A total of 169 responses were received from the survey, comprising 24% nephrologists (n=41), 36% cardiologists (n=61), and 40% emergency medicine physicians (n=67).

In terms of years of experience, the majority of nephrologists (61%) and approximately one-third of cardiologists (39%) had >10 years of experience. Conversely, an equal proportion of emergency medicine physicians (31%) had 5-10 years (31%) and >10 years (30%) of experience. Specifically, among nephrologists, 17% had 5-10 years, 15% had 2- <5 years, and 7% had <2 years of experience. For cardiologists, 21% had 5-10 years, 20% had 2- <5 years, and 18% had <2 years of experience. Among emergency medicine physicians, 27% had 2- <5 years, and 12% had <2 years of experience.

The majority of respondents (58%) were from Ministry of Health (MoH) hospitals, followed by 26% from private hospitals and 16% from university hospitals. Among nephrologists, 49% were from MoH hospitals, 34% from private hospitals, and 17% from university hospitals. Cardiologists were evenly split between MoH hospitals (44%) and private hospitals (44%), with a smaller percentage from university hospitals (11%). Most emergency medicine physicians (76%) were from MoH hospitals, with 19% from university hospitals and 4% from private hospitals.

The SC reviewed the votes and comments in detail. All statements achieved a level of agreement of 75% or higher among the participants (Appendix 1). Following the review, the 20 statements were further reduced and refined to 16 final statements (Table 1). The final statements were based on the latest available evidence. In cases where evidence was lacking, or recommendations were not feasible in the Malaysian context, a consensus was reached among the SC members. The finalised statements were designed to provide clear, evidence-based recommendations for managing acute and persistent hyperK⁺ in the Malaysian healthcare system. In addition, algorithms for managing acute (Figure 1) and persistent (Figure 2) hyperK⁺ were developed.

TARGET USER

These consensus statements are intended to guide those involved in the management of patients presenting with acute hyperK⁺, having persistent (chronic) hyperK⁺ or at risk of hyperK⁺ in primary, secondary and tertiary healthcare settings in the public and private sectors. It namely targets medical officers, specialists, trainee specialists and medical students.

SOURCE OF FUNDING

The development of these consensus statements was supported by an unrestricted grant from AstraZeneca Malaysia. AstraZeneca Malaysia played no role in the discussions and consensus process.

DECLARATION OF CONFLICT OF INTEREST

Dr Ching Chen Hua, Dr Mohd Rahal Bin Yusoff, Dr Shaik Farid Abdul Wahab, Dr Azmee bin Mohd Ghazi and Dr Siti Suhaila Hamzah declare no conflict of interest.

Dr Lim Soo Kun has received honorarium for presenting lectures and Advisory Boards from Astellas, AstraZeneca, Baxter, Boehringer Ingelheim, Bayer, Duopharma, Fresenius-Kabi, MSD, Novartis, Novo Nordisk, Roche, Sanofi and Taisho.

Dr Liew Houg Bang has received support for attending meetings and/or travel by AstraZeneca Sdn Bhd.

Dr Paranthaman Kaneson has received honoraria for presenting lectures from AstraZeneca Sdn Bhd.

Dr Prasad Menon has received honoraria for presenting lectures from AstraZeneca Sdn Bhd, Viatrix and ZP Therapeutics.

Dr Sunita Bavanandan has received honoraria for presenting lectures from Boehringer Ingelheim, Bayer and AstraZeneca Sdn Bhd. She also holds the following leadership roles: Chair of ISN OSEA Regional Board, Honorary Secretary Asian Pacific Society of Nephrology, Executive Committee Member KDIGO, and a member of the National Kidney Foundation Board of Directors.

Dr Tan Li Ping has received honorarium for presenting lectures and other education events from Sanofi, Boehringer Ingelheim, Novartis, Novo Nordisk, AstraZeneca, Fresenius, Duopharma and Bayer. He has also received payment for expert testimonies and participated in Advisory Boards for Bayer, Novo Nordisk, Fresenius and Boehringer Ingelheim. Additionally, he holds leadership roles in the Malaysian Society of Nephrology and the National Kidney Foundation. He holds multiple stock or stock options, but not in AstraZeneca.

Dr David Chew Soon Ping has received honorarium for presenting lectures from Johnson & Johnson and ZP Therapeutics, and participated in an Advisory Board for Otsuka. He was a committee member for the development of the 2023 Malaysian Clinical Practice Guidelines for Heart Failure.

ACKNOWLEDGMENT

The development committee would like to express their gratitude to the College of Emergency Physicians, the Malaysian Society of Nephrologists, and the National Heart Association of Malaysia councils for distributing the survey to their members, and all the respondents for participating in the survey and their comments and feedback on the consensus statements.


TABLE 1. CONSENSUS STATEMENTS ON THE MANAGEMENT OF ACUTE AND PERSISTENT HYPERKALAEMIA

Definition of hyperK⁺
1. The definitions of hyperK ⁺ are – mild: K ⁺ 5.5-5.9 mmol/L, moderate: K ⁺ 6.0-6.4 mmol/L and severe: K ⁺ ≥6.5 mmol/L.
Diagnosing and managing hyperK⁺ in emergency settings
2. It is essential to consider the entire clinical picture when diagnosing and treating patients with hyperK ⁺ because it can manifest with non-specific symptoms or be asymptomatic.
3. All patients presenting to the ED with serum K ⁺ >5.5 mmol/L should undergo an ECG.
4. Initiating treatment strategies in the ED is recommended for patients with serum K ⁺ >6.0 mmol/L regardless of ECG changes and in all patients with hyperkalaemic ECG changes.
5. Reassess serum K ⁺ levels, and monitor ECG, vital signs, and plasma glucose at a frequency appropriate to the clinical context until the serum K ⁺ improves to <5.5 mmol/L.
6. Patients who have responded to treatment (i.e., serum K ⁺ <5.5 mmol/L), are stable, with no indication for admission or risk of recurrent hyperK ⁺ may be discharged with preventive care and a follow-up appointment.
Preventing hyperK⁺ in the at-risk cardiorenal patients
7. Consider preventive strategies for patients at risk of hyperK ⁺ .
8. Consider using a K ⁺ binder for RAASi optimisation in patients with known hyperK ⁺ when basic preventive strategies fail.
Assessing risk and managing hyperK⁺ in cardiorenal patients
9. When RAASi are initiated, closer serum K ⁺ monitoring is recommended for high-risk patients (patients with CKD, HF, and/or DM).
10. In an individualised care plan, effectively managing hyperK ⁺ is recommended to optimise GDMT.
11. In cases of mild and moderate hyperK ⁺ , consider decreasing the RAASi dosage if mitigation strategies prove ineffective for normalising serum K ⁺ levels.
12. In cases of severe hyperK ⁺ , consider de-escalating or discontinuing RAASi when the risks outweigh the benefits of continuation.
K⁺-lowering therapies for hyperK⁺ in cardiorenal patients
13. Managing persistent hyperK ⁺ with long-term use of novel K ⁺ binders (SZC or patiromer) may enable cardiorenal patients to experience the proven benefits of guideline-recommended doses of RAASi therapy.
14. CPS is an option for short-term hyperK ⁺ management. Exercise caution when using CPS in the medium- or long-term due to concerns about GI side effects.
Collaborative care
15. In all cases presenting to the ED with moderate and severe hyperK ⁺ and are unresponsive to treatment, a physician consultation is recommended.
16. To optimise GDMT, and align medication prescriptions and adjustments relating to hyperK ⁺ , cross-specialty communication is essential.

CKD, chronic kidney disease; CPS, calcium polystyrene sulfonate; DM, diabetes mellitus; ECG, electrocardiogram; ED, emergency department; GDMT, guideline-directed medical treatment; GI, gastrointestinal.


gastrointestinal; HF, heart failure; hyperK⁺, hyperkalaemia; K⁺, potassium; RAASi, renin-angiotensin-aldosterone system inhibitors; SZC, sodium zirconium cyclosilicate.

FIGURE 1. EMERGENCY MANAGEMENT OF HYPERKALAEMIA (HYPERK⁺)



All K⁺ levels are in mmol/L. *Give 10 U soluble insulin + 50 ml Dextrose 50% (or equivalent strength) through a slow bolus. ABCDE, Airway, Breathing, Circulation, Disability, Exposure approach; CPS, calcium polystyrene sulfonate; ECG, electrocardiogram; ESKD, end-stage kidney disease; ICU, intensive care unit; IV, intravenous; K⁺, potassium; PO, orally; SZC, sodium zirconium cyclosilicate; TCA, to come again; TDS, three times daily. Adapted from The Renal Association UK. Clinical Practice Guidelines. 2020.¹

FIGURE 2. MANAGEMENT OF HYPERKALAEMIA (HYPERK⁺) IN THE OUTPATIENT SETTING



All K⁺ levels are in mmol/L. *RAAS-i = ACE-i/ARB, ARNI and MRA. ACE-i, angiotensin-converting enzyme inhibitor; AKI, acute kidney injury; ARB, angiotensin II receptor blocker; ARNI, angiotensin receptor-neprilysin inhibitor; CKD, chronic kidney disease; CPS, calcium polystyrene sulfonate; ECG, electrocardiogram; ED, emergency department; HF, heart failure; K⁺, potassium; MRA, mineralocorticoid receptor antagonist; NSAID, non-steroidal anti-inflammatory drug; PO, orally; TDS, three times daily; RAAS-i, renin-angiotensin-aldosterone system inhibitor; SZC, sodium zirconium cyclosilicate. Adapted from The Renal Association UK. Clinical Practice Guidelines. 2020.¹

1.0 INTRODUCTION

In healthy individuals, serum potassium (K^+) levels are tightly regulated and maintained between 3.5-5.0 mmol/L through a balance of intake and excretion.^{2,3}

- Studies have shown a U-shaped relationship between serum K^+ levels and mortality, with increased mortality observed when K^+ levels deviate from the normal range.^{3,4}
- In patients with heart failure (HF), chronic kidney disease (CKD), and diabetes mellitus (DM), a large retrospective study (n=911,698) found that all-cause mortality increases when serum K^+ levels fall outside the 4.0-5.0 mmol/L range.
- The study also observed that mortality rises with each 0.1 mmol/L increment above 5.0 mmol/L.⁴

Patients with DM, HF, and particularly CKD are at a significantly higher risk of developing hyperkalaemia (hyper K^+) than the general population.^{2,4,5}

- The use of renin-angiotensin-aldosterone system inhibitors (RAASi), comprising of angiotensin-converting enzyme inhibitors/angiotensin II receptor blockers (ACE-i/ARB), angiotensin receptor/neprilysin inhibitors (ARNI), and mineralocorticoid receptor antagonists (MRA), as a component of guidelines-directed medical therapy (GDMT)⁶⁻¹⁰ for HF, CKD with proteinuria, and DM with hypertension, further increases this risk.
 - In clinical trials involving renin-angiotensin system (RAS) blockers (ACEi/ARB and MRA), the rates of hyper K^+ events were reported to be 5.0% to 20.0%.¹¹⁻¹⁴
- Chronic (persistent) hyper K^+ , defined as a recurrent elevation in serum K^+ above normal limits, can lead to adverse cardiac effects and is common among patients with HF, CKD, and those on RAASi, resulting in medication changes in nearly 50% of affected patients.¹⁵

Even with the availability of effective novel K^+ binders like patiomer and sodium zirconium cyclosilicate (SZC) for long-term use, and the K^+ -lowering resin, calcium polystyrene sulfonate (CPS),¹⁶⁻¹⁸ concerns about the increased risk of hyper K^+ due to RAASi in these high-risk populations have hindered the optimisation of RAASi therapy.^{11,12,19,20}

- Sub-optimal RAASi dosing occurs although evidence shows that it worsens long-term mortality and hospitalisation rates in HF patients and increases the risk of cardiovascular (CV) events in CKD patients.^{11,12,21-26}

Despite this, ambiguity remains about the definition and thresholds of hyper K^+ and managing acute and persistent hyper K^+ in the cardiorenal population. These consensus statements aim to offer recommendations for managing acute and persistent hyper K^+ , incorporating the perspectives of emergency physicians, nephrologists, and cardiologists. These recommendations are based on the latest evidence, international guidelines, and best clinical practices within the Malaysian healthcare system.

2.0 Consensus Statements

1. Definition of hyperkalaemia

1.1. The definitions of hyperK⁺ are – mild: K⁺ 5.5-5.9 mmol/L, moderate: K⁺ 6.0-6.4 mmol/L and severe: K⁺ ≥6.5 mmol/L.

The Kidney Disease: Improving Global Outcomes (KDIGO) suggests the classification of hyperK⁺ severity should consider complementing the absolute serum K⁺ level with the presence or absence of hyperkalaemic-based electrocardiogram (ECG) changes.²⁷

- However, the KDIGO recommendations are specifically for patients with CKD, and these patients have a lower risk of mortality associated with corresponding hyperK⁺ severity than other populations.²⁸
 - Patients with no CKD had a significantly higher risk of mortality at serum K⁺ 5.5-5.9 mmol/L (Odds ratio [OR] 10.32) than at serum K⁺ <5.5 mmol/L.
 - The risks tripled (OR 31.64) when serum K⁺ levels were ≥6.0 mmol/L.

Additionally, it is crucial to ensure that the high serum K⁺ levels are not due to **pseudohyperK⁺**.

- PseudohyperK⁺ (factitious, artefactual and spurious hyperK⁺) is a falsely elevated serum K⁺ despite a normal in vivo K⁺ concentration.²⁹
- Some causes of pseudohyperK⁺ include:³⁰
 - fist clenching during blood draw
 - lysed samples (due to tourniquet time >1 minute, mechanical trauma, using a pneumatic tube without cushioning, use of fine gauge needles, and duration of blood sample storage)
 - thrombocytosis
 - leucocytosis

Therefore, based on the evidence and current best practice, **this consensus guidance defines hyperK⁺** without including ECG changes.

The definitions of hyperK⁺ are:

- i. Mild hyperK⁺: serum K⁺ 5.5-5.9 mmol/L
- ii. Moderate hyperK⁺: serum K⁺ 6.0-6.4 mmol/L
- iii. Severe hyperK⁺: serum K⁺ ≥6.5 mmol/L

2. Diagnosing and managing hyperkalaemia in emergency settings

ECG changes arise when the balance between intracellular and extracellular K⁺ is disrupted, typically with serum K⁺ >5.5 mmol/L.

- This imbalance increases cell membrane excitability, which can lead to muscle weakness and, in severe cases, cardiac arrest during diastole.^{31,32}
- However, while ECG changes are a useful indicator of deviations in serum K⁺ levels, **hyperK⁺ often presents without noticeable signs and symptoms** and is frequently detected incidentally through laboratory tests conducted for other medical issues.³³

Life-threatening cardiac dysrhythmias generally occur at serum K⁺ levels >6.5 mmol/L, though they can manifest at lower levels in some patients.

- Conversely, those with persistent hyperK⁺ may not exhibit ECG changes and can remain asymptomatic even at higher K⁺ levels (see Statement 2.2).^{27,33}
- Additionally, it is important to consider that **symptomatic manifestations of hyperK⁺ depend not only on the absolute serum K⁺ level but also on the rate at which it increases.**³³

Assessing hyperK⁺ accurately and initiating emergency treatment, especially in the emergency department (ED), is challenging due to uncertainties about the rate of serum K⁺ increase and distinguishing acute from persistent hyperK⁺.

- Comprehensive patient history and clinical assessment are essential for appropriate treatment.
- ECG changes that could lead to lethal arrhythmias underscore the need for timely management and ongoing reassessment after initiating acute treatment.

Although current hyperK⁺ guidelines provide recommendations for monitoring serum K⁺ levels,^{1,30,34,35} they do not fully address essential parameters such as cardiac monitoring and monitoring for hypoglycaemia. Generally, patients whose serum K⁺ declines to ≤5.5 mmol/L may be discharged from the ED.³⁶

The algorithm for the emergency management of hyperK is presented in Figure 1.

2.1. It is essential to consider the entire clinical picture when diagnosing and treating patients with hyperK⁺ because it can manifest with non-specific symptoms or be asymptomatic.

Practice points:

2.1.1 Assess patients for hyperK⁺ even in the absence of symptoms, as life-threatening cardiac arrhythmias may occur at lower serum K⁺ levels and in patients with persistent hyperK⁺.

2.1.2 Be aware of non-specific hyperkalaemic symptoms for timely diagnosis and management.

Patients with hyperK⁺ may present with non-specific symptoms like:^{33,37,38}

- fatigue
- muscle weakness or pain
- paraesthesia
- flaccid paralysis
- depressed deep tendon reflexes
- palpitations
- syncope
- dyspnoea
- vomiting or nausea

When assessing the full clinical presentation, identifying risk factors of hyperK⁺ is essential. These include,

- a history of concurrent comorbidities (CKD, diabetic kidney disease [DKD], HF)
- medication history
- presenting complaints and other investigative findings indicating other causes of acute hyperK⁺, e.g., infections, uncontrolled hyperglycaemia, and acute kidney injury (AKI).

2.2. All patients presenting to the ED with serum K⁺>5.5 mmol/L must undergo an ECG.

Practice points:

2.2.1 Although ECG sensitivity for mild hyperK⁺ is low, increased serum K⁺ can induce detectable ECG changes.

Changes in the ECG can occur even with mild hyperK⁺, such as tall, peaked, narrow-based T waves in the V2-V4 leads and fascicular blocks.^{31,32,37}

- Progressively increasing serum K⁺ induces further ECG changes such as different types of heart blocks, decreasing P-wave amplitude that can be followed by the disappearance of the P-wave, prolonged QRS complex, ventricular dysrhythmias, “sine waves”, and finally asystole, in severe hyperK⁺.^{31,32,37}

The severity of hyperK⁺ and ECG changes can be variable, and the **absence of an ECG abnormality does not negate the presence of hyperK⁺**.^{32,37} Despite this, **assessing ECG is an essential screening tool** to assess the severity of hyperK⁺ rapidly.³⁹

2.3. Initiating treatment strategies in the ED is recommended for patients with serum K⁺>6.0 mmol/L regardless of ECG changes and in all patients with hyperkalaemic ECG changes.

Practice points:

2.3.1 Accurately differentiating between acute and persistent hyperK⁺ is essential to guide appropriate emergency treatments.

2.3.2. Promptly initiate emergency management strategies for hyperK⁺ based on the serum K⁺ level to prevent ECG changes and potential lethal arrhythmia.

2.3.3. To manage hyperK⁺ in the emergency setting, treat reversible causes, reduce membrane excitability (e.g., with intravenous [IV] calcium gluconate), and start measures for lowering serum K⁺.

Managing hyperK⁺ in the emergency setting consists of four basic steps that are well-established and widely practiced (Figure 1):^{33,34,40-45}

1. Determine and treat any reversible cause of the hyperK⁺.
2. Take measures to reduce the cardiac membrane excitability with IV calcium salts, like calcium gluconate.
3. Initiate therapeutic measures to reduce serum K⁺ levels rapidly:
 - a. Initial treatment should be with IV 50% dextrose and insulin. This combination is useful for patients with relative contraindications to β-2

- agonists and those with severe hyperK⁺ (serum K⁺ ≥6.0 mmol/L or associated ECG changes).
- b. Subsequently, salbutamol can be administered by nebuliser or IV route.
 - c. Loop or thiazide diuretics may be considered for non-oliguric and volume-overloaded patients. IV frusemide 40 mg can be given every 12 hours or as a continuous infusion for patients with volume overload and preserved kidney function.^{33,34}
 - d. K⁺-binders may be considered for patients with renal insufficiency.^{33,34} However, SZC cannot be used as a stand-alone therapy in life-threatening acute hyperK⁺ and should be used in addition to the standard of care.⁴⁶
 - e. Sodium bicarbonate is useful in patients with severe acidosis.
4. If the serum K⁺ remains ≥6.5 mmol/L or pathological ECG changes persist, the renal team should be contacted to arrange for urgent dialysis, as haemodialysis is the definitive measure to reduce serum K⁺.

2.4. Reassess serum K⁺ levels, and monitor ECG, vital signs, and plasma glucose at a frequency appropriate to the clinical context until the serum K⁺ improves to <5.5 mmol/L.

Practice points:

2.4.1 Monitoring ECG continuously at and during emergency management of hyperK⁺ is advisable.

2.4.2 Monitor other clinical parameters at appropriate intervals on a case-to-case basis.

2.4.3 Monitor glucose levels at regular intervals if the patient is given insulin during the emergency treatment of hyperK⁺.

2.4.4 Monitor serum K⁺ levels as early 2-4 hours after initiating K⁺-lowering agents and then reassess to determine the frequency of monitoring.

The frequency and duration of monitoring depend on:³⁰

- the severity of the hyperK⁺ and its manifestation
- the type of K⁺ lowering drugs administered
- the likelihood of rebound hyperK⁺
- the patient's overall clinical context and response to treatment

Monitoring parameters	Methods
ECG	<ul style="list-style-type: none"> • Continuous ECG and/or cardiac monitoring is suggested for all patients treated for acute hyperK⁺ with serum K⁺ >6.0 mmol/L.³⁰
Other vital signs	<ul style="list-style-type: none"> • Other vital signs, such as blood pressure and oxygen saturation, should also be monitored.³⁰
Hypoglycaemia	<ul style="list-style-type: none"> • Can occur as early as 30 minutes to up to 3.5 hours^{47,48} post-insulin-glucose administration. • May persist up to 6 hours.¹

	<ul style="list-style-type: none"> Monitor glucose frequently within the first hour, followed by appropriate intervals, depending on the insulin and glucose dose administered, pre-treatment glucose levels⁴⁹ and other patient-specific risk factors.⁵⁰
Serum K⁺	<p><i>K⁺-shifting agents</i></p> <ul style="list-style-type: none"> Structured monitoring is typically aligned with the action times of the K⁺-shifting agents, which begin within 30-60 minutes. As these medications do not remove K⁺ from circulation, rebound hyperK⁺ can potentially occur after 2-3 hours.³⁰ Re-evaluation of serum K⁺ can be performed as early as 2-4 hours after administration of agents like insulin-glucose infusion and salbutamol. <p><i>K⁺-binders</i></p> <ul style="list-style-type: none"> Using K⁺ binders may help mitigate the rebound effect, though their onset of action varies, <ul style="list-style-type: none"> older binders like CPS can take hours to days to exert their effect,⁵¹ newer binders like SZC act within an hour and patiromer within 4-7 hours.⁵² Serum K⁺ levels can be measured at 1-2 hours post-K⁺ binder administration,¹ followed by regular monitoring at appropriate intervals until the patient is discharged from the ED.

2.5. Patients who have responded to treatment (i.e., serum K⁺ <5.5 mmol/L), are stable, with no indication for admission or risk of recurrent hyperK⁺ may be discharged with preventive care and a follow-up appointment.

Practice points:

2.5.1 Healthcare providers may discharge patients who have a repeat serum K⁺ <5.5 mmol/L, resolved ECG changes, and stable vital parameters, with K⁺ binders and close follow-up.

2.5.2 Hemodynamic instability, new or persistent ECG changes, and new onset hyperK⁺ require admission.

The recent consensus statement from the American College of Emergency Physicians³⁵ recommends **admission for further evaluation** for patients who,

- do not achieve haemodynamic stability
- continue developing new ECG abnormalities
- have recalcitrant serum K⁺ levels
- have no history of hyperK⁺

Consider discharging patients from the ED if they fulfil the following criteria:

- History of persistent hyperK⁺ and the cause for the present acute episode has been resolved.

2. Achieving haemodynamic stability and feeling well.
3. Has easy access and avenues for close follow-up.
4. Agrees to discharge after being counselled about the risks and benefits of discharge.

Before discharge, the physician should:

1. Review the patient's current medications and diet.
2. Advise them to withhold any K⁺ supplements and to consume a low K⁺ diet.
3. Ensure a K⁺ binder has been prescribed with the correct dose and duration.
4. Provide a referral letter to the nearest healthcare facility for a repeat serum K⁺ on completion of the K⁺ binder.
5. Ensure that the precipitating factor has been identified and managed accordingly.

3. Preventing hyperK⁺ in the at-risk cardiorenal patients

K⁺ homeostasis is primarily maintained by the kidneys, which excrete K⁺.⁵³

- Impairment in this process can lead to hyperK⁺.
- Several medications can disrupt the K⁺ balance by either causing a transcellular shift of K⁺ or by reducing renal excretion.⁵⁴
- Conditions like CKD, HF, DM and hypertension, especially when treated with RAASi, elevate the risk of persistent hyperK⁺, making patients 2-3 times more likely to develop hyperK⁺.^{52,53,55,56}
- Additional risk factors include using other drugs like β-blockers, non-steroid anti-inflammatory drugs (NSAIDs), and trimethoprim/sulfamethoxazole, advanced age, and high K⁺ intake.^{52,55,57} Hence, preventing hyperK⁺ in these at-risk patients is crucial to optimising guideline-directed medical treatment (GDMT) use for better patient outcomes.

3.1. Consider preventive strategies for patients at risk of hyperK⁺

Practice points:

3.1.1 Consider regular blood monitoring for patients at risk of hyperK⁺ and introduce preventive strategies when serum K⁺ is ≥ 5.0 mmol/L.

3.1.2 RAASi should be carefully dosed, and serum K⁺ levels should be monitored closely after initiation and up-titration.

3.1.3 Using diuretics can mitigate hyperK⁺ in patients with volume overload.

3.1.4 Address modifiable risk factors such as avoiding medications and drug combinations that may cause hyperK⁺, avoiding constipation, and correcting metabolic acidosis.

3.1.5 Dietary modification limiting K⁺-rich foods, withdrawing K⁺-containing supplements and educating patients about hidden K⁺ in processed foods are also preventive strategies.

Prescribe hyperK⁺-inducing drugs with caution – The following drugs should be prescribed with caution for individuals who are at high risk of persistent hyperK⁺.⁵⁴

1. Drugs inhibiting the RAAS: These drugs interfere with K⁺ homeostasis, causing a reduction in K⁺ excretion, leading to hyperK⁺. They include RAASi, NSAIDs,

calcineurin inhibitors (e.g., cyclosporine and tacrolimus), heparin and its derivatives, other aldosterone antagonists, K⁺-sparing diuretics, trimethoprim and pentamidine.

2. Medications altering transmembrane K⁺ movement: They include amino acids, β-blockers, suxamethonium and mannitol.
3. K⁺-containing agents: Like salt substitutes and alternatives, penicillin G and stored blood products.

Diet modification – Guidelines recommend that patients at risk of hyperK⁺ should reduce their K⁺ intake in their daily diet.^{1,34,56}

- However, there are practical and psychological barriers to adhering to a low K⁺ diet, and limited evidence of the association between dietary K⁺, serum K⁺, and hyperK⁺.⁵⁸
- The latest National Kidney Foundation’s Kidney Disease Outcomes Quality Initiative (KDQI) guidelines⁵⁹ recommend other K⁺ lowering strategies to effectively prevent hyperK⁺ and specify that adjusting dietary K⁺ intake for adults with CKD stages 3-5D or post-transplantation may be reasonable.
- KDIGO (2024) guidelines⁶⁰ recommend educating patients with CKD about dietary modifications (including K⁺ intake) based on their individual needs, the severity of their CKD and other comorbid conditions.
- Patients with CKD or at risk of hyperK⁺ should also be educated about **sources of hidden K⁺ in processed foods**.⁶¹
 - K⁺ additives in processed foods are used as preservatives, stabilisers, and in flavouring agents and sweeteners.
 - Examples of processed foods with hidden K⁺ include dried fruits, peanut butter, meat derivatives, tomato-based sauces,⁶¹ sweetened and alcoholic beverages and packaged snacks.

Using diuretics – Diuretics, particularly loop diuretics, enhance K⁺ excretion through urine and are commonly prescribed to prevent an increase of serum K⁺ and control overload simultaneously.^{27,56}

- Diuretics should be dosed based on the kidney function because their K⁺-lowering efficacy reduces as kidney impairment advances.
- However, caution must be used by regularly assessing the patient’s kidney function and fluid status, as diuretics may cause AKI and electrolyte imbalance, including hypokalaemia.
- In addition, diuretics might not be ideal for long-term use to control serum K⁺ unless there are other reasons for it, such as extracellular volume overload.⁵⁶

Addressing metabolic acidosis – If there is concurrent metabolic acidosis (HCO₃ <22 mmol/L), oral sodium bicarbonate (3-5 g/day) can be considered.

- However, prolonged use of sodium bicarbonate may increase the sodium load, resulting in volume overload. In this circumstance, sodium bicarbonate should be discontinued.^{27,56}

3.2. Consider using a K⁺ binder for RAASi optimisation in patients with known hyperK⁺ when basic preventive strategies fail.

Practice points:

3.2.1 Use optimal doses of RAASi for managing CKD because it reduces albuminuria, increases the rate of albuminuria normalisation and slows CKD progression.

3.2.2 Optimise RAASi for managing HF as it reduces hospitalisations for HF, CV mortality and all-cause mortality.

3.2.3 Consider initiating K⁺ binders to facilitate RAASi optimisation in patients with hyperK⁺.

Full-dose irbesartan has been shown to significantly reduce the progression to macroalbuminuria over two years, lowering the incidence from 14.9% to 5.2% while also increasing the regression to normoalbuminuria when compared to placebo.⁶²

- Additionally, optimal antiproteinuric doses of benazepril and losartan have been found to reduce the risk of doubling serum creatinine, end-stage kidney disease (ESKD), or death by 51% and 53%, respectively, compared to conventional dosages in non-diabetic CKD patients.⁶³
- Higher doses of lisinopril have been demonstrated to reduce all-cause mortality or hospitalisation for HF compared to lower doses.⁶⁴
- Eplerenone has also been shown to reduce CV-related mortality, all-cause mortality, and HF-related hospitalisations.⁶⁵
- Maximally tolerated doses of sacubitril/valsartan have been found to reduce the risk of CV death, first hospitalisation for HF, and all-cause mortality when compared to enalapril.⁶⁶

Novel K⁺ binders like SZC and patiomer can help optimise dosing of RAASi in patients at higher risk of hyperkalaemia.

- SZC has been shown to effectively correct and maintain normal serum K⁺ levels in patients with hyperkalaemia, with 78% achieving normal levels after a 24-72-hour correction phase.⁶⁷
- Over a 12-month maintenance period, up to three-quarters of patients were able to maintain their RAASi dose.
- Patiomer has been found to lower the risk of hyperK⁺ (>5.5 mmol/L) and reduce the need for MRA dose adjustments in patients with HF with reduced ejection fraction and a current history of RAASi-related hyperK⁺.⁶⁸

4. Assessing risk and managing hyperkalaemia in cardiorenal patients

A population-based analysis revealed that 28% of CKD and 39% of HF patients experienced at least one episode of hyperK⁺.⁵⁶

- A history of hyperK⁺ episodes increases the likelihood of recurrence, underscoring the importance of ongoing vigilance.⁵⁶
- Since individuals at risk for persistent hyperK⁺ may not exhibit symptoms, regular monitoring is essential for early detection and management.

Before considering dose de-escalation or withdrawal, guidelines from the European Society of Cardiology (ESC),¹⁰ American Heart Association/American College of Cardiology (AHA/ACC),⁹ and the KDIGO guidelines^{6,8} recommend,

- optimising RAASi to the highest tolerated dose,
- close monitoring for hyperK⁺, and
- addressing hyperK⁺.

The algorithm for managing hyperK⁺ in the outpatient setting (persistent hyperK⁺) is presented in Figure 2.

4.1. When RAASi are initiated, closer serum K⁺ monitoring is recommended for high-risk patients (patients with CKD, HF, and/or DM).

Practice points:

4.1.1 RAASi elevates hyperK⁺ risk in patients with comorbidities like CKD, HF, and DM, necessitating closer serum K⁺ monitoring.

Patients with CKD	Have a proportionally increasing risk of developing hyperK ⁺ with more advanced levels of kidney impairment. ⁶⁹
Patients with DM	Have an increased risk of hyperK ⁺ due to the associated hyporeninemic hypoaldosteronism. ^{13,70}
Patients with HF	Are at risk due to multiple factors, including medications used to manage this condition. ^{13,70}

These risks are additive; hence, an individual with more than one of these conditions (e.g., CKD with HF) would be at a higher risk of hyperK⁺ than an individual with only one of these conditions.⁷¹ For details on initiating and monitoring RAASi, see Statements 4.3 and 4.4.

4.2. In an individualised care plan, effectively managing hyperK⁺ is recommended to optimise GDMT.

Practice points:

4.2.1 Patients with HF on GDMT require regular laboratory monitoring for hyperK⁺ and preventive actions against AKI.

4.2.2 To manage hyperK⁺ effectively without interrupting GDMT, prioritise managing all modifiable risk factors.


When patients at risk of hyperK⁺ are prescribed GDMTs or other non-disease modifying agents, attention must be given to,

- patient education
- repeat laboratory monitoring
- preventing AKI, which can precipitate hyperK⁺

Shared-decision making is essential in personalising care, and patients should be adequately counselled regarding the risks and benefits of GDMT.^{12,22,25}

If hyperK⁺ occurs, effective management involves addressing contributing factors like, diet, use of salt substitutes, acidosis and medications (e.g. K⁺-sparing diuretics and K⁺ supplements) before reducing or discontinuing GDMT. (see Statements 3.1-3.3) Figure 3 is an example of an individualised care plan to manage hyperK⁺ while optimising GDMT effectively.⁶

Figure 3. Monitoring serum creatinine and K⁺ during ACEi or ARB treatment with dose adjustment and monitoring of side effects.



ACE, angiotensin converting enzyme inhibitor; AKI, acute kidney injury; ARB, angiotensin receptor blockers; hyperK⁺, hyperkalaemia; K⁺, potassium; NSAIDs, non-steroidal anti-inflammatory drugs. Source: KDIGO 2022 Clinical Practice Guidelines for Diabetes Management in Chronic Kidney Disease.⁶ Creative Commons Attribution CC BY-NC-ND-4.0.

4.3. In cases of mild and moderate hyperK⁺, consider decreasing the RAASi dosage if mitigation strategies prove ineffective.

Practice points:

4.3.1 Ensure that all other strategies to mitigate hyperK⁺ have been optimised (see Statements 3.1, 3.2)

4.3.2 Consider a K⁺ binder, if available, to avoid RAASi dose reduction. (see Statements 5.1 and 5.2)

4.3.3 Consider reducing RAASi dose and monitor serum K⁺ level if long-term use of K⁺ binder is not feasible.

4.4. In cases of severe hyperK⁺, consider discontinuing RAASi when the risks outweigh the benefits of continuation.

Practice points:

4.4.1 Ensure all other strategies to mitigate hyperK⁺ have been optimised (see Statements 3.1, 3.2)

4.4.2 To determine if RAASi can be restarted, temporarily discontinue RAASi and evaluate renal function before deciding on a treatment approach.

4.4.3 Consider a K⁺ binder, if available, to facilitate reinitiating RAASi. (see Statements 5.1 and 5.2)

The initiation and target doses of RAASi for treating patients with HF can be found within the ESC¹⁰ and AHA/ACC⁹ guidelines, while the KDIGO guidelines^{6,8} recommend optimising RAASi up to the maximal tolerated dose for managing blood pressure.

- Before considering dose reduction or temporary discontinuation of RAASi, all other strategies to mitigate hyperK⁺, including using oral K⁺ binders, should have been optimised.

Patients with serum K⁺ >5.0 mmol/L⁷²

- RAASi should be started at a low dose and up-titrated cautiously.
- Using novel K⁺ binders may be considered to help optimise RAASi utilisation in patients who are either not started on GDMT or have their GDMT not optimised.

Patients with serum K⁺ >6.5 mmol/L⁷²

- If clinically indicated, RAASi should be temporarily discontinued, and emergency treatment should be initiated to resolve hyperK⁺.
- Consider novel K⁺ binders upon RAASi re-initiation once serum K⁺ is normalised to <5.5 mmol/L post-acute treatment.

- Serum K⁺ level and kidney function must be evaluated with every dose adjustment made in patients with serum K⁺ >5.0 mmol/L.
- Frequency of approaching serum K⁺ monitoring has been suggested for ACEi, ARBs and MRAs in Table 3.

Table 3. Frequency of serum K⁺ monitoring for ACEi, ARB and MRAs.

- At treatment initiation
- 2-4 weeks after initiation but earlier if the patient has kidney impairment with an interval depending on the severity of impairment

- At regular intervals, thereafter, based on the patient's individual risk

5. Potassium lowering therapies for hyperkalaemia in cardiorenal patients

Discontinuation of RAASi treatment following a single hyperK⁺ episode in cardiorenal patients is common despite guidelines recommending continuation.⁷³

- Only a small proportion of patients (up to 15%) had RAASi reinstated within six months, with up to 37% having their RAASi dose reduced by more than 25%.⁷³
- Reduced RAASi treatment after a hyperK⁺ episode increased all-cause hospital inpatient days by up to 18.2 days within six months.⁷⁴
- It has been reported that hyperK⁺-related RAASi discontinuation or down-titration was associated with a higher risk of cardiorenal events compared to maintained or up-titrated RAASi dosing in patients with CKD or HF.

5.1. Managing persistent hyperK⁺ with long-term use of novel K⁺ binders (SZC or patiomer) may enable cardiorenal patients to experience the proven benefits of guideline-recommended doses of RAASi therapy.

Practice points:

5.1.1 Persistent hyperK⁺ is one of the most important causes of RAASi dosage de-escalation or discontinuation.

5.1.2 Consider using novel K⁺ binders to optimise the use of RAASi as they have demonstrated long-term efficacy and safety.

Novel K⁺ binders (patiomer and SZC) have demonstrated efficacy in reducing serum K⁺ compared to placebo, and safety for long-term use up to 12 months in cardiorenal patients.¹⁶⁻¹⁸

- This potentially supports GDMT optimisation and reduces the risk of worse outcomes.
- Generally, the trials involving the novel K⁺ binders involved patients with CKD, DM or HF with hyperK⁺.
- The duration of these Phase III trials, including the randomised controlled phase, ranged from 12 days to 27 weeks.
- A few of the studies with an open-labelled maintenance phase ran up to 12 months.²⁷ The important trials on novel K⁺ binders, including patiomer and SZC, are summarised in Tables 4 and 5.

Table 4. Patiromer Phase III trials

No	Trial	Design	Population	Duration	Findings
1.	OPAL-HK ⁷⁵	Multicentre phase 1: single group, single arm. Phase 2: randomised, single-blind, placebo-controlled withdrawal phase.	Phase 1: N=243 Phase 2: N=107 eGFR <60 (91%), RAASi (100%), DM (57%), HTN (97%), hyper ^{K+}	Phase 1: 4 weeks Phase 2: 8 weeks	Mean serum K ⁺ reduction in phase 1 was -1.01 mmol/L (p<0.001). Between group difference in serum K ⁺ during the withdrawal phase was 0.72 mmol/L (95% CI 0.46–0.99, p<0.001). 60% of the placebo group compared with 15% of the patiromer group had a serum K ⁺ reading ≥5.5 mmol/L.
2.	DIAMOND ⁶⁸	Multicentre Phase 1: open label, run in. Phase 2: randomised triple-blind placebo-controlled withdrawal phase.	Phase 1: N=1195 Phase 2: N=878 RAASi (100%), MRA (100%), HTN (91%), HF (100%), hyper ^{K+}	Phase 1: up to 12 weeks Phase 2: 13 to 43 weeks (median: 27 weeks)	Serum K ⁺ was 0.10 mmol/L higher in the placebo group (p<0.001). Serum K ⁺ >5.5 mmol/L and discontinuation or dose reduction of MRA was less frequent in the patiromer group (HR 0.63 and 0.62, with p=0.006 for both).

CI, confidence interval; DM, diabetes mellitus; eGFR, estimated glomerular filtration rate; HR, hazard ratio; HTN, hypertension; hyper^{K+}, hyperkalaemia; K⁺, potassium; MRA, mineralocorticoid antagonists; RAASi, renin angiotensin-aldosterone system inhibitors.

Table 5. SZC Phase III trials

No	Trial	Design	Population	Duration	Findings
1.	ZS-003 ¹⁷	Phase 1: multicentre, double-blind correction phase. Phase 2: randomised controlled, multicentre, double-blind maintenance phase.	Phase 1: N=753 Phase 2: N=543 eGFR ≤60 (74.5%), RAASi (67%), DM (60%), hyperK ⁺	Phase 1: 48 hours Phase 2: 12 days	In the correction phase, SZC reduced serum K ⁺ by 0.30% per hour while placebo reduced serum K ⁺ by 0.09% per hour (p<0.001). During the maintenance phase, SZC 5 g and 10 g significantly decreased mean serum K ⁺ at days 9 and 15 (5.11 mmol/L vs 4.62 mmol/L and 5.11 mmol/L vs 4.60 mmol/L, respectively; p<0.001 for both) but not at day 21 (5.14 mmol/L vs 4.96 mmol/L, p=0.221). There were no significant between group changes in arrhythmias.
2.	HARMONIZE ⁷⁶	Phase 1: multicentre, open-label correction phase. Phase 2: randomised controlled, multicentre, double-blind maintenance phase.	Phase 1: N=251 Phase 2: N=237 eGFR ≤60 (69%), RAASi (72%), DM (68%), hyperK ⁺	Phase 1: 48 hours Phase 2: 28 days	By the end of Phase 1, the mean reduction in serum K ⁺ was - 1.1 mmol/L (p<0.001). In Phase 2, normal serum K ⁺ levels were obtained more often in the SZC arms (71%, p=0.01; 76%, p=0.002; 85%, p<0.001 for 5 g, 10 g and 15 g, respectively). The differences remained significant in the HF subgroup.

3.	HARMONIZE Open-label extension ¹⁸	Multicentre, open-label, single-arm, maintenance phase extension.	N=121 eGFR ≤60 (74%), RAASi (69%), DM (67%), hyperK ⁺	≥11 months	Mean serum K ⁺ ≤5.1 mmol/L was achieved by 76.6% to 87.5% of participants, and mean serum K ⁺ ≤5.5 mmol/L was obtained by all participants.
4.	ZS-005 ⁶⁷	Phase 1: multicentre, open-label correction phase. Phase 2: multicentre, open-label, maintenance phase.	Phase 1: N=751 Phase 2: N=746 eGFR ≤60 (73%), RAASi (64%), DM (63%), HTN (82%), hyperK ⁺	Phase 1: 24–72 hours Phase 2: up to 12 months	At completion of the correction phase, 78% of the patients had a normal serum K ⁺ . During the maintenance phase, 74% of participants maintained their RAASi dose.
5.	Kashihara et al. ⁷⁷	RCT, multicentre, double-blind (Phase II/III)	N=103 eGFR ≤60 (97%), RAASi (78%), DM (60%)	48 hours	At 48 hours, the proportion of patients with normal serum K ⁺ was higher in the SZC 5 g and 10 g arms versus placebo (85.3% and 91.7% vs 15.2%, respectively; p<0.0001).
6.	Kashihara et al. ¹⁶	Multicentre, open-label, single arm correction and maintenance phase.	N=150 eGFR ≤60 (93%), RAASi (71%), DM (58%), hyperK ⁺	Phase 1: 1–3 days Phase 2: 1 year	99% of the patients in the correction phase had normal serum K ⁺ after 72 h. ≥65.5% of patients had normokalaemia throughout.
7.	DIALIZE ⁷⁸	RCT, multicentre, double-blind.	N=196 mHD (100%), pre-dialysis hyperK ⁺	8 weeks	More participants in the SZC arm had pre-dialysis serum K ⁺ levels between 4 and 5 mmol/L in at least 3/4 long interdialytic intervals without requiring rescue therapy versus placebo (41.2% vs 1.0%; p<0.001). Rescue therapy for hyperK ⁺ was similar in both arms.

8.	ENERGIZE ⁴⁶	RCT, multicentre, double-blind.	N=70 Emergency room, hyperK ⁺	10 hours	There was no statistically significant reduction in serum K ⁺ between SZC and placebo at 10 hours.
----	------------------------	---------------------------------	---	----------	---

CI, confidence interval; DM, diabetes mellitus; eGFR, estimated glomerular filtration rate; HF, heart failure; HR, hazard ratio; HTN, hypertension; hyperK⁺, hyperkalaemia; K⁺, potassium; mHD, maintenance haemodialysis; MRA, mineralocorticoid antagonists; RAASi, renin-angiotensin-aldosterone system inhibitors; RCT, randomised controlled trial; SZC, sodium zirconium cyclosilicate.

5.2. CPS is an option for short-term hyperK⁺ management. Exercise caution when using CPS in the medium- or long-term due to concerns about gastrointestinal (GI) side effects.

Practice points:

5.2.1 The K⁺ binder, CPS, is an option for the short-term management of hyperK⁺.

5.2.2 CPS is administered orally. If the oral route is not feasible or a rapid onset of action is needed, the rectal route can be used.

5.2.3 CPS use in the medium- and long-term should be done with caution as it may cause bowel necrosis, GI intolerance, hypercalcaemia, and hypomagnesemia.

CPS is a K⁺-lowering resin that exchanges calcium for K⁺.

- It acts in the colon and can be administered orally or rectally if the oral route is not an option or rapid onset is needed.^{79,80}
- CPS has demonstrated efficacy in reducing serum K⁺ in hyperK⁺ among patients with CKD (Table 5).^{79,81,82}
- Even so, an extensive systematic review to determine the efficacy and safety of these resins revealed that there is still a lack of high-quality evidence for using them to treat hyperK⁺.⁸⁰
- The most common and major side effect of CPS is constipation.⁸³⁻⁸⁵
 - There are some case reports of patients developing intestinal necrosis, resulting from a combined administration of CPS and sorbitol.⁸³⁻⁸⁵
 - Since there is a small risk of serious intestinal complications, patients prescribed CPS should be closely observed.
 - The product information for CPS recommends that it should be discontinued if there are any abnormal GI side effects, such as severe constipation, prolonged abdominal pain, and vomiting.⁷⁹
 - Data for CPS long-term efficacy is limited. A retrospective study (n=247 with CKD stages 2-5) showed that long-term CPS (mean CPS dose 8.0 ± 3.6 g/day and mean duration 5.6 ± 8.7 months) may be effective and safe for controlling mild hyperK⁺ in patients with CKD.⁸⁵

Table 5. CPS clinical trials

No	Trial	Design	Population	Duration	Findings
1.	Wang et al., 2018	Prospective, randomised, crossover clinical trial with a 1-week washout period.	58 haemodialysis patients with hyperK ⁺ (≥5.5 mol/L).	3-week CPS (3 x 5 g/day) or a blank control, with a 1-week washout period.	Compared with the control group, CPS treatment significantly reduced serum K ⁺ levels (p <0.01). More patients in the CPS group had lower serum K ⁺ levels than the safety level of <5.5 mmol/L (32% for control vs. 61% for CPS, p <0.01).
2.	Wang et al., 2023	Prospective, open, randomised, controlled, single-centre clinical	107 stage 3–5 non-dialysis CKD patients with hyperK ⁺ group A (15 g/day) or group B (30 g/day).	1 week	After 3 days of treatment, the serum K ⁺ levels in groups A and B had decreased by 0.68 ± 0.46 and 0.75 ± 0.43 mmol/L, respectively.

		observational study.			After 7 days, the serum K ⁺ levels in groups A and B had decreased by 0.64 ± 0.37 and 0.94 ± 0.49mmol/L, respectively.
--	--	----------------------	--	--	---

CKD, chronic kidney disease; CPS, calcium polystyrene sulfonate; K⁺, potassium.

6. Collaborative care

The use of RAASi in managing closely linked diseases requires close cross-speciality treatment approaches to ensure better patient outcomes.^{70,86-89}

- The risk of hyperK⁺ with RAASi is less than 2.0% if used as monotherapy, and increases to approximately 5.0% in dual RAASi usage, and up to 10.0% if combined dual therapy is used in patients with HF and CKD.⁹⁰
- It is worth noting that in the real-world setting, the incidence of hyperK⁺ can be as high as 50.0%.⁹¹⁻⁹⁴
- This is most probably due to starting RAASi in unselected or poorly selected patients who did not receive recommended serum K⁺ and kidney function monitoring.^{93,94}

6.1. In all cases presenting to the ED with moderate and severe hyperK⁺ and are unresponsive to treatment, a physician consultation is recommended.

Practice points:

6.1.1 Refer patients unresponsive to emergency hyperK⁺ management to nephrologists/physicians.

6.1.2 Ensure referral/follow-up of patients to work up the cause of acute hyperK⁺ and prevent its recurrence.

6.1.3 Refer patients with persistent ECG abnormalities to the relevant specialities.

Patients with moderate and severe hyperK⁺, with or without ECG changes, who do not respond to the initial acute treatment should be referred to the nephrologist for dialysis.

- A K⁺ binder may also be considered a short-term measure to control the serum K⁺ levels.
- These patients should also be referred to a physician to monitor for rebound hyperK⁺ after the initial reduction.
- Further workup is needed to determine the cause of hyperK⁺ and to prevent recurrence.
- A cardiology review is required to ensure cardiac stability if persistent ECG abnormalities are present.¹

The patient should be monitored in an intensive care setting (e.g. coronary care unit, high dependency unit or the intensive care unit), particularly if they are:

- Anuric with ESKD.
- Severely dehydrated.
- Exhibiting ECG abnormalities with haemodynamic instability.
- Exhibiting an altered conscious level.

6.2. To optimise GDMT, and align medication prescriptions and adjustments relating to hyperK⁺, cross-specialty communication is essential.

Practice points:

6.2.1 Clear channels of communication between cardiology, nephrology, endocrinology, internal medicine, and dietetics is essential to optimise patient care and manage comorbidities associated with hyperK⁺ and RAAS-i therapy.

Clear communication regarding therapy in patients with HF and CKD should be established. In some cases, close collaboration with dietitians and pharmacists is required to evaluate the patient's diet, use of supplements, salt substitutes and nutraceuticals, and other drugs that can increase the serum K⁺. Quality initiative programs should be encouraged to monitor the kidney function and K⁺ levels frequently.

Conclusion

The effective management of hyperK⁺ is essential for improving patient outcomes, especially in those with cardiorenal conditions. Optimal management should aim to reduce ED visits and unplanned hospitalisations due to hyperK and its associated complications while allowing optimisation of GDMT. This multidisciplinary set of consensus statements provides comprehensive guidelines based on the latest evidence and best practices specific to the context of the Malaysian healthcare system.

By standardising the approach to diagnosing and treating both acute and persistent hyperK⁺, these recommendations aim to reduce complications, optimise the use of GDMTs like RAASi, and ultimately improve the quality of care for at-risk patients. The availability of novel potassium binders, such as patiomer and SZC, provides clinicians with effective long-term options that enable the safe and optimal use of RAASi in high-risk populations. Through these efforts, healthcare providers can better prevent and manage hyperK⁺, minimising its impact on patients and the healthcare system.

Appendix: Agreement for the 20 consensus statements

Statement	Agreement
1. The definitions of hyperK ⁺ are – mild: K ⁺ >5.5 mmol/L (5.6-6.0), moderate: K ⁺ >6.0 mmol/L (6.1-6.5) and severe: K ⁺ >6.5 mmol/L.	93%
2. HyperK ⁺ can manifest with non-specific symptoms or be asymptomatic. Hence, it is essential to consider the entire clinical picture when diagnosing and treating patients with hyperK ⁺ .	96%
3. Baseline ECG should be done for all patients presenting to the ED with K ⁺ >6 mmol/L and is recommended for K ⁺ >5.5 mmol/L.	91%
4. Acute treatment strategies should be initiated in patients with K ⁺ >6 mmol/L regardless of baseline ECG findings and in all patients with hyperkalaemic-ECG changes with K ⁺ >5.5 mmol/L.	90%
5. Reassessment of K ⁺ levels, and ECG, vital signs and plasma glucose monitoring are recommended at a frequency appropriate to the clinical context until the K ⁺ level is normalised.	96%
6. Post-acute treatment, patients with a normalised K ⁺ level who are stable and do not have any other cause for admission or risk of recurrent hyperK ⁺ may be discharged with preventive care and follow-up.	92%
7. K ⁺ -binding agents, loop-diuretics and dialysis are the means to remove K ⁺ from the body.	98%
8. Reducing ED visits and unplanned hospitalisations due to complications associated with hyperK ⁺ should be a goal of good optimal management.	98%
9. Patients with CKD, HF, and/or DM and patients treated RAASi are at increased risk of hyperK ⁺ .	97%
10. Effective management of hyperK ⁺ should be an integral part of an individualised care plan to optimise GDMT that improves morbidity, mortality and patient outcomes.	97%
11. In mild and moderate hyperK ⁺ , the de-escalation or discontinuation of GDMTs such as RAASi should only be done as a last resort.	94%
12. In severe hyperK ⁺ , de-escalation or discontinuation of GDMT can be considered when the risks outweigh the benefits of continuation.	98%
13. For high-risk patients currently not hyperkalaemic, preventative measures, such as avoidance of certain non-disease modifying agents and foods, should be considered.	98%
14. If basic preventive measures have not been successful, patients with a known history of hyperK ⁺ prohibiting the optimisation of RAASi may benefit from using a novel K ⁺ binder to facilitate RAASi optimisation.	94%
15. Patients at risk of hyperK ⁺ should be monitored closely with a strategy in place to manage K ⁺ levels effectively.	97%
16. The long-term use of novel potassium binders (SZC or patiromer) for managing persistent hyperK ⁺ may enable guideline-recommended doses of RAASi therapy and allow cardiorenal patients to experience their proven benefits.	98%
17. The use of SPS should be avoided due to concerns about severe GI toxicity, low compliance due to poor palatability.	87%

18. CPS should be used with caution in the medium- or long-term as it may cause GI side effects, including bowel necrosis.	97%
19. Nephrology consultation is recommended for all patients presenting to the ED with moderate and severe hyperK ⁺ unresponsive to typical emergency treatment.	94%
20. Cross-speciality communication is essential to align medication prescriptions and adjustments, in addition to optimising GDMTs.	98%

CKD, chronic kidney disease; CPS, calcium polystyrene sulfonate; DM, diabetes mellitus; ECG, electrocardiogram; ED, emergency department; GDMT, guideline-directed medical treatment; GI, gastrointestinal; HF, heart failure; hyperK⁺, hyperkalaemia; K⁺, potassium; RAASi, renin-angiotensin-aldosterone system inhibitors; SPS, sodium polystyrene sulfonate; SZC, sodium zirconium cyclosilicate.

References

1. The Renal Association. Clinical practice guidelines treatment of acute hyperkalaemia in adults. 2020. Available: https://ukkidney.org/sites/renal.org/files/RENAL%20ASSOCIATION%20HYPERKALAEMIA%20GUIDELINE%20-%20JULY%202022%20V2_0.pdf. Accessed March 2024.
2. Fitch K, Woolley JM, Engel T, Blumen H. The Clinical and Economic Burden of Hyperkalemia on Medicare and Commercial Payers. *Am Health Drug Benefits*. Jun 2017;10(4):202-210.
3. Kovesdy CP. Updates in hyperkalemia: Outcomes and therapeutic strategies. *Rev Endocr Metab Disord*. Mar 2017;18(1):41-47. doi:10.1007/s11154-016-9384-x
4. Collins AJ, Pitt B, Reaven N, et al. Association of Serum Potassium with All-Cause Mortality in Patients with and without Heart Failure, Chronic Kidney Disease, and/or Diabetes. *Am J Nephrol*. 2017;46(3):213-221. doi:10.1159/000479802
5. Nilsson E, Gasparini A, Årnlöv J, et al. Incidence and determinants of hyperkalemia and hypokalemia in a large healthcare system. *Int J Cardiol*. Oct 15 2017;245:277-284. doi:10.1016/j.ijcard.2017.07.035
6. Kidney Disease Improving Global Outcomes (KDIGO). Clinical practice guideline for diabetes management in chronic kidney disease 2022. Available: <https://kdigo.org/wp-content/uploads/2022/10/KDIGO-2022-Clinical-Practice-Guideline-for-Diabetes-Management-in-CKD.pdf>. Accessed March 2024.
7. Ministry of Health Malaysia. Clinical Practice Guidelines Management of Heart Failure 2023 (5th ed). Available at: <https://www.malaysianheart.org/publication/clinical-practice-guidelines/p/clinical-practice-guidelines-management-of-heart-failure-2023-5th-edition>.
8. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. Mar 2021;99(3s):S1-s87. doi:10.1016/j.kint.2020.11.003
9. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. May 3 2022;79(17):e263-e421. doi:10.1016/j.jacc.2021.12.012
10. McDonagh TA, Metra M, Adamo M, et al. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J*. Sep 21 2021;42(36):3599-3726. doi:10.1093/eurheartj/ehab368
11. Beusekamp JC, Tromp J, Cleland JGF, et al. Hyperkalemia and Treatment With RAAS Inhibitors During Acute Heart Failure Hospitalizations and Their Association With Mortality. *JACC Heart Fail*. Nov 2019;7(11):970-979. doi:10.1016/j.jchf.2019.07.010
12. Epstein M, Reaven NL, Funk SE, McGaughey KJ, Oestreich N, Knispel J. Evaluation of the treatment gap between clinical guidelines and the utilization of renin-angiotensin-aldosterone system inhibitors. *Am J Manag Care*. Sep 2015;21(11 Suppl):S212-20.
13. Ismail U, Sidhu K, Zieroth S. Hyperkalaemia in Heart Failure. *Card Fail Rev*. Mar 2021;7:e10. doi:10.15420/cfr.2020.29
14. Leon SJ, Tangri N. Balancing Hyperkalemia Risks with Clinical Benefits of Renin-Angiotensin-Aldosterone Inhibitors/Mineralocorticoid Receptor Antagonists Blockade: It's Apples and Oranges. *Kidney360*. Aug 25 2022;3(8):1442-1444. doi:10.34067/kid.0000952022
15. Riccio E, Capuano I, Buonanno P, et al. RAAS Inhibitor Prescription and Hyperkalemia Event in Patients With Chronic Kidney Disease: A Single-Center Retrospective Study. *Front Cardiovasc Med*. 2022;9:824095. doi:10.3389/fcvm.2022.824095
16. Kashihara N, Yamasaki Y, Osonoi T, et al. A phase 3 multicenter open-label maintenance study to investigate the long-term safety of sodium zirconium cyclosilicate in Japanese subjects with hyperkalemia. *Clin Exp Nephrol*. Feb 2021;25(2):140-149. doi:10.1007/s10157-020-01972-y

17. Packham DK, Rasmussen HS, Lavin PT, et al. Sodium zirconium cyclosilicate in hyperkalemia. *N Engl J Med*. Jan 15 2015;372(3):222-31. doi:10.1056/NEJMoa1411487
18. Roger SD, Spinowitz BS, Lerma EV, et al. Efficacy and Safety of Sodium Zirconium Cyclosilicate for Treatment of Hyperkalemia: An 11-Month Open-Label Extension of HARMONIZE. *Am J Nephrol*. 2019;50(6):473-480. doi:10.1159/000504078
19. Fonseca C, Brito D, Branco P, Frazão JM, Silva-Cardoso J, Bettencourt P. Hyperkalemia and management of renin-angiotensin-aldosterone system inhibitors in chronic heart failure with reduced ejection fraction: A systematic review. *Rev Port Cardiol (Engl Ed)*. Sep 2020;39(9):517-541. doi:10.1016/j.repc.2020.03.015
20. Henrysson J, Thunström E, Chen X, Fu M, Basic C. Hyperkalaemia as a cause of undertreatment with mineralocorticoid receptor antagonists in heart failure. *ESC Heart Fail*. Feb 2023;10(1):66-79. doi:10.1002/ehf2.14137
21. Fu EL, Evans M, Clase CM, et al. Stopping Renin-Angiotensin System Inhibitors in Patients with Advanced CKD and Risk of Adverse Outcomes: A Nationwide Study. *J Am Soc Nephrol*. Feb 2021;32(2):424-435. doi:10.1681/asn.2020050682
22. Linde C, Bakhai A, Furuland H, et al. Real-World Associations of Renin-Angiotensin-Aldosterone System Inhibitor Dose, Hyperkalemia, and Adverse Clinical Outcomes in a Cohort of Patients With New-Onset Chronic Kidney Disease or Heart Failure in the United Kingdom. *J Am Heart Assoc*. Nov 19 2019;8(22):e012655. doi:10.1161/jaha.119.012655
23. Muhlestein JB, Kammerer J, Bair TL, et al. Frequency and clinical impact of hyperkalaemia within a large, modern, real-world heart failure population. *ESC Heart Fail*. Feb 2021;8(1):691-696. doi:10.1002/ehf2.13164
24. O'Connor CM, Abraham WT, Albert NM, et al. Predictors of mortality after discharge in patients hospitalized with heart failure: an analysis from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF). *Am Heart J*. Oct 2008;156(4):662-73. doi:10.1016/j.ahj.2008.04.030
25. Singhanian G, Ejaz AA, McCullough PA, et al. Continuation of Chronic Heart Failure Therapies During Heart Failure Hospitalization - a Review. *Rev Cardiovasc Med*. Sep 30 2019;20(3):111-120. doi:10.31083/j.rcm.2019.03.562
26. Walther CP, Winkelmayr WC, Richardson PA, Virani SS, Navaneethan SD. Renin-angiotensin system blocker discontinuation and adverse outcomes in chronic kidney disease. *Nephrol Dial Transplant*. Sep 27 2021;36(10):1893-1899. doi:10.1093/ndt/gfaa300
27. Larivée NL, Michaud JB, More KM, Wilson JA, Tennankore KK. Hyperkalemia: Prevalence, Predictors and Emerging Treatments. *Cardiol Ther*. Mar 2023;12(1):35-63. doi:10.1007/s40119-022-00289-z
28. Einhorn LM, Zhan M, Hsu VD, et al. The frequency of hyperkalemia and its significance in chronic kidney disease. *Arch Intern Med*. Jun 22 2009;169(12):1156-62. doi:10.1001/archinternmed.2009.132
29. Kapoor AK, Ravi A, Twomey PJ. Investigation of outpatients referred to a chemical pathologist with potential pseudohyperkalaemia. *J Clin Pathol*. Oct 2009;62(10):920-3. doi:10.1136/jcp.2008.062141
30. Lindner G, Burdmann EA, Clase CM, et al. Acute hyperkalemia in the emergency department: a summary from a Kidney Disease: Improving Global Outcomes conference. *Eur J Emerg Med*. Oct 2020;27(5):329-337. doi:10.1097/mej.0000000000000691
31. Teymouri N, Mesbah S, Navabian SMH, et al. ECG frequency changes in potassium disorders: a narrative review. *Am J Cardiovasc Dis*. 2022;12(3):112-124.
32. Van Mieghem C, Sabbe M, Knockaert D. The clinical value of the ECG in noncardiac conditions. *Chest*. Apr 2004;125(4):1561-76. doi:10.1378/chest.125.4.1561
33. Simon LV, Hashmi MF, Farrell MW. Hyperkalemia. [Updated 2023 Sep 4]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470284/>.

34. Clase CM, Carrero JJ, Ellison DH, et al. Potassium homeostasis and management of dyskalemia in kidney diseases: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference. *Kidney Int.* Jan 2020;97(1):42-61. doi:10.1016/j.kint.2019.09.018
35. Rafique Z, Peacock F, Armstead T, et al. Hyperkalemia management in the emergency department: An expert panel consensus. *J Am Coll Emerg Physicians Open.* Oct 2021;2(5):e12572. doi:10.1002/emp2.12572
36. Singer AJ, Thode HC, Jr., Peacock WF. Rapid correction of hyperkalemia is associated with reduced mortality in ED patients. *Am J Emerg Med.* Nov 2020;38(11):2361-2364. doi:10.1016/j.ajem.2019.12.012
37. Campese VM, Adenuga G. Electrophysiological and clinical consequences of hyperkalemia. *Kidney Int Suppl (2011).* Apr 2016;6(1):16-19. doi:10.1016/j.kisu.2016.01.003
38. Te Dorsthorst RPM, Hendrikse J, Vervoorn MT, van Weperen VYH, van der Heyden MAG. Review of case reports on hyperkalemia induced by dietary intake: not restricted to chronic kidney disease patients. *Eur J Clin Nutr.* Jan 2019;73(1):38-45. doi:10.1038/s41430-018-0154-6
39. Diercks DB, Shumaik GM, Harrigan RA, Brady WJ, Chan TC. Electrocardiographic manifestations: electrolyte abnormalities. *J Emerg Med.* Aug 2004;27(2):153-60. doi:10.1016/j.jemermed.2004.04.006
40. Guidelines and Audit Implementation Network (GAIN). Guidelines for the treatment of hyperkalaemia in adults. 2014. Available: <https://www.rqia.org.uk/RQIA/files/6f/6f51b366-f8bf-44de-a630-6967d5353a87.pdf>. Accessed March 2024.
41. Dépret F, Peacock WF, Liu KD, Rafique Z, Rossignol P, Legrand M. Management of hyperkalemia in the acutely ill patient. *Ann Intensive Care.* Feb 28 2019;9(1):32. doi:10.1186/s13613-019-0509-8
42. Martinez-Vea A, Bardají A, Garcia C, Oliver JA. Severe hyperkalemia with minimal electrocardiographic manifestations: a report of seven cases. *J Electrocardiol.* Jan 1999;32(1):45-9. doi:10.1016/s0022-0736(99)90020-1
43. Sims DB, Sperling LS. ST-Segment Elevation Resulting From Hyperkalemia. *Circulation.* 2005;111(19):e295-e296. doi:doi:10.1161/01.CIR.0000165127.41028.D1
44. Webster A, Brady W, Morris F. Recognising signs of danger: ECG changes resulting from an abnormal serum potassium concentration. *Emerg Med J.* Jan 2002;19(1):74-7. doi:10.1136/emj.19.1.74
45. Wong R, Banker R, Aronowitz P. Electrocardiographic changes of severe hyperkalemia. *J Hosp Med.* Apr 2011;6(4):240. doi:10.1002/jhm.656
46. Peacock WF, Rafique Z, Vishnevskiy K, et al. Emergency Potassium Normalization Treatment Including Sodium Zirconium Cyclosilicate: A Phase II, Randomized, Double-blind, Placebo-controlled Study (ENERGIZE). *Acad Emerg Med.* Jun 2020;27(6):475-486. doi:10.1111/acem.13954
47. UK Renal Association. Clinical Practice Guidelines: Treatment of Acute Hyperkalemia in Adults. Available: <http://www.renal.org/guidelines/joint-guidelines/treatment-of-acutehyperkalemia-in-adults-sthash.GI1GDFeb.dpbs>. Accessed March 2024.
48. Schafers S, Naunheim R, Vijayan A, Tobin G. Incidence of hypoglycemia following insulin-based acute stabilization of hyperkalemia treatment. *J Hosp Med.* Mar 2012;7(3):239-42. doi:10.1002/jhm.977
49. Chothia MY, Humphrey T, Schoonees A, Chikte UME, Davids MR. Hypoglycaemia due to insulin therapy for the management of hyperkalaemia in hospitalised adults: A scoping review. *PLoS One.* 2022;17(5):e0268395. doi:10.1371/journal.pone.0268395
50. Boughton CK, Dixon D, Goble E, et al. Preventing Hypoglycemia Following Treatment of Hyperkalemia in Hospitalized Patients. *J Hosp Med.* May 2019;14(5):284-287. doi:10.12788/jhm.3145

51. Synapse. What is calcium polystyrene sulfonate used for. Available at: <https://synapse.patsnap.com/article/what-is-calcium-polystyrene-sulfonate-used-for>.
52. Palmer BF, Carrero JJ, Clegg DJ, et al. Clinical Management of Hyperkalemia. *Mayo Clin Proc.* Mar 2021;96(3):744-762. doi:10.1016/j.mayocp.2020.06.014
53. Di Lullo L, Ronco C, Granata A, et al. Chronic Hyperkalemia in Cardiorenal Patients: Risk Factors, Diagnosis, and New Treatment Options. *Cardiorenal Med.* 2019;9(1):8-21. doi:10.1159/000493395
54. Ben Salem C, Badreddine A, Fathallah N, Slim R, Hmouda H. Drug-induced hyperkalemia. *Drug Saf.* Sep 2014;37(9):677-92. doi:10.1007/s40264-014-0196-1
55. Wang AY. Optimally managing hyperkalemia in patients with cardiorenal syndrome. *Nephrol Dial Transplant.* Dec 1 2019;34(Suppl 3):iii36-iii44. doi:10.1093/ndt/gfz225
56. Weinstein J, Girard LP, Lepage S, McKelvie RS, Tennankore K. Prevention and management of hyperkalemia in patients treated with renin-angiotensin-aldosterone system inhibitors. *Cmaj.* Dec 6 2021;193(48):E1836-e1841. doi:10.1503/cmaj.210831
57. Neuen BL, Oshima M, Agarwal R, et al. Sodium-Glucose Cotransporter 2 Inhibitors and Risk of Hyperkalemia in People With Type 2 Diabetes: A Meta-Analysis of Individual Participant Data From Randomized, Controlled Trials. *Circulation.* May 10 2022;145(19):1460-1470. doi:10.1161/circulationaha.121.057736
58. Ramos CI, González-Ortiz A, Espinosa-Cuevas A, Avesani CM, Carrero JJ, Cuppari L. Does dietary potassium intake associate with hyperkalemia in patients with chronic kidney disease? *Nephrol Dial Transplant.* Nov 9 2021;36(11):2049-2057. doi:10.1093/ndt/gfaa232
59. Ikizler TA, Burrowes JD, Byham-Gray LD, et al. KDOQI Clinical Practice Guideline for Nutrition in CKD: 2020 Update. *Am J Kidney Dis.* Sep 2020;76(3 Suppl 1):S1-s107. doi:10.1053/j.ajkd.2020.05.006
60. KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int.* Apr 2024;105(4s):S117-s314. doi:10.1016/j.kint.2023.10.018
61. Martínez-Pineda M, Vercet A, Yagüe-Ruiz C. Are Food Additives a Really Problematic Hidden Source of Potassium for Chronic Kidney Disease Patients? *Nutrients.* Oct 12 2021;13(10)doi:10.3390/nu13103569
62. Parving HH, Lehnert H, Bröchner-Mortensen J, Gomis R, Andersen S, Arner P. The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med.* Sep 20 2001;345(12):870-8. doi:10.1056/NEJMoa011489
63. Hou FF, Xie D, Zhang X, et al. Renoprotection of Optimal Antiproteinuric Doses (ROAD) Study: a randomized controlled study of benazepril and losartan in chronic renal insufficiency. *J Am Soc Nephrol.* Jun 2007;18(6):1889-98. doi:10.1681/asn.2006121372
64. Packer M, Poole-Wilson PA, Armstrong PW, et al. Comparative effects of low and high doses of the angiotensin-converting enzyme inhibitor, lisinopril, on morbidity and mortality in chronic heart failure. ATLAS Study Group. *Circulation.* Dec 7 1999;100(23):2312-8. doi:10.1161/01.cir.100.23.2312
65. Zannad F, McMurray JJ, Krum H, et al. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med.* Jan 6 2011;364(1):11-21. doi:10.1056/NEJMoa1009492
66. Bhatt AS, Vaduganathan M, Claggett BL, et al. Effect of sacubitril/valsartan vs. enalapril on changes in heart failure therapies over time: the PARADIGM-HF trial. *Eur J Heart Fail.* Sep 2021;23(9):1518-1524. doi:10.1002/ejhf.2259
67. Spinowitz BS, Fishbane S, Pergola PE, et al. Sodium Zirconium Cyclosilicate among Individuals with Hyperkalemia: A 12-Month Phase 3 Study. *Clin J Am Soc Nephrol.* Jun 7 2019;14(6):798-809. doi:10.2215/cjn.12651018
68. Butler J, Anker SD, Lund LH, et al. Patiromer for the management of hyperkalemia in heart failure with reduced ejection fraction: the DIAMOND trial. *Eur Heart J.* Nov 1 2022;43(41):4362-4373. doi:10.1093/eurheartj/ehac401

69. Thomsen RW, Nicolaisen SK, Hasvold P, et al. Elevated potassium levels in patients with chronic kidney disease: occurrence, risk factors and clinical outcomes-a Danish population-based cohort study. *Nephrol Dial Transplant*. Sep 1 2018;33(9):1610-1620. doi:10.1093/ndt/gfx312
70. Bianchi S, Aucella F, De Nicola L, Genovesi S, Paoletti E, Regolisti G. Management of hyperkalemia in patients with kidney disease: a position paper endorsed by the Italian Society of Nephrology. *J Nephrol*. Aug 2019;32(4):499-516. doi:10.1007/s40620-019-00617-y
71. Latts LM, Reaven NL, Funk SE, McGaughey KJ, Adamson RT. Hyperkalemia Is Highly Prevalent In Patients With Cardiorenal Comorbidities Compared To Patients Without These Comorbidities. *Value in Health*. 2015;18(3):A135. doi:10.1016/j.jval.2015.03.786
72. Rosano GMC, Tamargo J, Kjeldsen KP, et al. Expert consensus document on the management of hyperkalaemia in patients with cardiovascular disease treated with renin angiotensin aldosterone system inhibitors: coordinated by the Working Group on Cardiovascular Pharmacotherapy of the European Society of Cardiology. *Eur Heart J Cardiovasc Pharmacother*. Jul 1 2018;4(3):180-188. doi:10.1093/ehjcvp/pvy015
73. Kanda E, Pollack Jr C, Rastogi A, et al. #3303 Suboptimal extents of RAASi re-initiation after discontinuation following hyperkalemia: an observational study of cardiorenal patients in the US and Japan. *Nephrology Dialysis Transplantation*. 2023;38(Supplement_1)doi:10.1093/ndt/gfad063c_3303
74. Svensson ME, Murohara T, Sundström J, et al. #4406 Increase in hospitalised days after hyperkalemia-related reduction in RAASi use: an observational study on cardiorenal patients in Sweden and Japan. *Nephrology Dialysis Transplantation*. 2023;38(Supplement_1)doi:10.1093/ndt/gfad063c_4406
75. Pitt B, Bakris GL, Bushinsky DA, et al. Effect of patiromer on reducing serum potassium and preventing recurrent hyperkalaemia in patients with heart failure and chronic kidney disease on RAAS inhibitors. *Eur J Heart Fail*. Oct 2015;17(10):1057-65. doi:10.1002/ejhf.402
76. Kosiborod M, Rasmussen HS, Lavin P, et al. Effect of sodium zirconium cyclosilicate on potassium lowering for 28 days among outpatients with hyperkalemia: the HARMONIZE randomized clinical trial. *Jama*. Dec 3 2014;312(21):2223-33. doi:10.1001/jama.2014.15688
77. Kashihara N, Nishio T, Osonoi T, et al. Correction of serum potassium with sodium zirconium cyclosilicate in Japanese patients with hyperkalemia: a randomized, dose-response, phase 2/3 study. *Clin Exp Nephrol*. Dec 2020;24(12):1144-1153. doi:10.1007/s10157-020-01937-1
78. Fishbane S, Ford M, Fukagawa M, et al. A Phase 3b, Randomized, Double-Blind, Placebo-Controlled Study of Sodium Zirconium Cyclosilicate for Reducing the Incidence of Predialysis Hyperkalemia. *J Am Soc Nephrol*. Sep 2019;30(9):1723-1733. doi:10.1681/asn.2019050450
79. Kalimate Package Insert. Available: <https://quest3plus.bpfk.gov.my/pmo2/index.php>. Available March 2024.
80. Wong SWS, Zhang G, Norman P, Welihinda H, Wijeratne DT. Polysulfonate Resins in Hyperkalemia: A Systematic Review. *Can J Kidney Health Dis*. 2020;7:2054358120965838. doi:10.1177/2054358120965838
81. Wang J, Lv MM, Zach O, et al. Calcium-Polystyrene Sulfonate Decreases Inter-Dialytic Hyperkalemia in Patients Undergoing Maintenance Hemodialysis: A Prospective, Randomized, Crossover Study. *Ther Apher Dial*. Dec 2018;22(6):609-616. doi:10.1111/1744-9987.12723
82. Wang X, Chen D, Song X, Wang J, Zhang H. Efficacy and safety of calcium polystyrene sulfonate in patients with hyperkalemia and stage 3-5 non-dialysis chronic kidney disease: a single-center randomized controlled trial. *J Int Med Res*. Apr 2023;51(4):3000605231167516. doi:10.1177/03000605231167516

83. Joo M, Bae WK, Kim NH, Han SR. Colonic mucosal necrosis following administration of calcium polystyrene sulfonate (Kalimate) in a uremic patient. *J Korean Med Sci*. Dec 2009;24(6):1207-11. doi:10.3346/jkms.2009.24.6.1207
84. Takeuchi N, Nomura Y, Meda T, Iida M, Ohtsuka A, Naba K. Development of Colonic Perforation during Calcium Polystyrene Sulfonate Administration: A Case Report. *Case Rep Med*. 2013;2013:102614. doi:10.1155/2013/102614
85. Yu MY, Yeo JH, Park JS, Lee CH, Kim GH. Long-term efficacy of oral calcium polystyrene sulfonate for hyperkalemia in CKD patients. *PLoS One*. 2017;12(3):e0173542. doi:10.1371/journal.pone.0173542
86. De Nicola L, Ferraro PM, Montagnani A, Pontremoli R, Dentali F, Sesti G. Recommendations for the management of hyperkalemia in patients receiving renin-angiotensin-aldosterone system inhibitors. *Intern Emerg Med*. Sep 29 2023;doi:10.1007/s11739-023-03427-0
87. Essa H, Walker L, Mohee K, et al. Multispecialty multidisciplinary input into comorbidities along with treatment optimisation in heart failure reduces hospitalisation and clinic attendance. *Open Heart*. 2022;9(2):e001979. doi:10.1136/openhrt-2022-001979
88. Ponikowski P, Voors AA, Anker SD, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. Jul 14 2016;37(27):2129-2200. doi:10.1093/eurheartj/ehw128
89. Rydén L, Grant PJ, Anker SD, et al. ESC guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD - summary. *Diab Vasc Dis Res*. May 2014;11(3):133-73. doi:10.1177/1479164114525548
90. Vardeny O, Wu DH, Desai A, et al. Influence of baseline and worsening renal function on efficacy of spironolactone in patients With severe heart failure: insights from RALES (Randomized Aldactone Evaluation Study). *J Am Coll Cardiol*. Nov 13 2012;60(20):2082-9. doi:10.1016/j.jacc.2012.07.048
91. Cooper LB, Hammill BG, Peterson ED, et al. Consistency of Laboratory Monitoring During Initiation of Mineralocorticoid Receptor Antagonist Therapy in Patients With Heart Failure. *Jama*. Nov 10 2015;314(18):1973-5. doi:10.1001/jama.2015.11904
92. Palmer BF. Managing hyperkalemia caused by inhibitors of the renin-angiotensin-aldosterone system. *N Engl J Med*. Aug 5 2004;351(6):585-92. doi:10.1056/NEJMra035279
93. Raebel MA. Hyperkalemia associated with use of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers. *Cardiovasc Ther*. Jun 2012;30(3):e156-66. doi:10.1111/j.1755-5922.2010.00258.x
94. Shah KB, Rao K, Sawyer R, Gottlieb SS. The adequacy of laboratory monitoring in patients treated with spironolactone for congestive heart failure. *J Am Coll Cardiol*. Sep 6 2005;46(5):845-9. doi:10.1016/j.jacc.2005.06.010